

$\frac{7}{10}$

Chapter 06

Mostafa Touny

October 3, 2023

Contents

Problems	2
1	2
2	2
4	2
8	2
11	3
12	3
14	3
21	3
22	3
24	3
34	4

Problems

1 $\phi: \mathbb{Z} \rightarrow 2\mathbb{Z}$

$\phi(n) = 2n$. If $2a = 2b$ then $a = b$. For each $2k$ we have $\phi(k) = 2k$. Observe $\phi(ab) = 2(a+b) = 2a + 2b = \phi(a)\phi(b)$, Following by usual properties of integers.

0.5
0.5

2

We Follow the same proof approach of *Example 15* (page 130). Let $\phi \in \text{Aut}(Z)$ be arbitrary. Then by the usual properties of integers and isomorphisms, $\phi(k) = \phi(1+1+\dots+1) = \phi(1) + \dots + \phi(1) = k \cdot \phi(1)$. But by definition $\phi(1) = c$ for some integer c . Therefore $\phi(k) = kc$. In other words, $\text{Aut}(Z) = \{\phi \mid \exists c, \forall k \phi(k) = kc\}$

Completely wrong! But the grade was for the approach. $\text{Aut}(Z) = \{\epsilon, -\epsilon\}$

ϕ
2
c needs to be either 1 or -1 for ϕ to be surjective!

Cayley table of $U(8)$:

	1	3	5	7
1	1	3	5	7
3	3	1	7	5
5	5	7	1	3
7	7	5	3	1

Cayley table of $U(10)$:

	1	3	7	9
1	1	3	7	9
3	3	9	1	7
7	7	1	9	3
9	9	7	3	1

Recall from *theorem 6.2* (page 126), Any ϕ maps the identity to the identity of the other group.

In $U(8)$ we have $3 \cdot 3 = 1$. Then $\phi(3 \cdot 3) = \phi(3) \cdot \phi(3) = \phi(1) = 1$. The only non-identity element in $U(10)$ satisfying that is 9. Hence $\phi(3) = 9$.

Similarly $5 \cdot 5 = 1$. Then we must have some $a \in U(10)$ such that $a \cdot a = 1$ where $a \notin \{1, 9\}$. Contradiction.

0.5
0.5

8

Injective. Given $\log_{10} a = \log_{10} b$, we get $10^{\log_{10} a} = 10^{\log_{10} b}$, and $a = b$.

Surjective. Given $x \in \mathcal{R}$, take $a = 10^x \in \mathcal{R}^+$. Then $\log_{10} a = \log_{10} 10^x = x$.

Group Operation. Observe $\phi(ab) = \log_{10} ab = \log_{10} a + \log_{10} b = \phi(a) + \phi(b)$.

0.5
0.5

11

Observe $\phi(a^3b^{-2}) = \phi(a^3) + \phi(b^{-2}) = [\phi(a)]^3 + [\phi(b)]^{-2} = (\bar{a})^3 + (\bar{b})^{-2}$. We used *theorem 6.2 (2)*.

0.8 $3\bar{a} - 2\bar{b}$

12

(\rightarrow). For any $a, b \in G$, We have:

0.5 shouldn't start with inverses without justification.
 $\alpha(a^{-1}b^{-1}) = \alpha(a^{-1})\alpha(b^{-1})$
 $(a^{-1}b^{-1})^{-1} = a b$
 $ba = ab$

(\leftarrow). Symmetrically, If we have $b^{-1}a^{-1} = a^{-1}b^{-1}$, Then $\alpha(ab) = \alpha(a)\alpha(b)$. Bijection is clear by properties of inverses. *Should clarify!*

1

14

By *theorem 6.5* (page 131), $Aut(Z_3) \approx U(3)$ and $Aut(Z_4) \approx U(4)$, so $Aut(Z_3) \approx Aut(Z_4)$ by the *transitivity* of isomorphism. But $Z_3 \not\approx Z_4$ as the two groups have different orders, so no bijection exists.

0.5

21

Clearly groups H and K are isomorphic to S_4 . By transitivity $H \approx K$.

Construct an isomorphism!

0.5
2

22

For every $c = 2, 3, 4, \dots$, Consider the subset $H_c = \{ck \mid k \in Z\}$. It is a subgroup, As it has the identity $c(0)$, inverses $c(-k)$, and closed $ck_1 + ck_2 = c(k_1 + k_2)$.

It remains to show those subgroups are distinct. For any c_1 and c_2 where $c_1 < c_2$ we have $c_1(1) \in H_{c_1}$ but $c_1(1) \notin H_{c_2}$. Therefore $H_{c_1} \neq H_{c_2}$.

1

24

We use *theorem 3.2* (page 63). If $\phi(a) = a$ then $\phi(a^{-1}) = (\phi(a))^{-1} = a^{-1}$. Also, If $\phi(a) = a$ and $\phi(b) = b$ then $\phi(ab) = \phi(a)\phi(b) = ab$.

0.375
0.5

It is nonempty because any aut. fixes the identity

34

Let K be a subgroup of G . We use *theorem 3.2* (page 63).

Inverse. For any $\phi(k) \in \phi(K)$, $(\phi(k))^{-1} = \phi(k^{-1})$. But $k^{-1} \in K$, So $\phi(k^{-1}) \in \phi(K)$.

for any $\bar{k} \in \phi(K)$

\bar{k} has a preimage in K by def. of $\phi(K)$.

Closed. For $\phi(k_1), \phi(k_2) \in \phi(K)$, We have $\phi(k_1)\phi(k_2) = \phi(k_1k_2)$. But $k_1k_2 \in K$, So $\phi(k_1k_2) \in \phi(K)$.

Q.E.D.