

$$\frac{9}{10}$$

## Chapter 08

Mostafa Touny

November 3, 2023

### Contents

<b>Problems</b>	<b>2</b>
4 . . . . .	2
5 . . . . .	2
6 . . . . .	2
15 . . . . .	2
17 . . . . .	3
21 . . . . .	3
23 . . . . .	3
35 . . . . .	4
46 . . . . .	4
48 . . . . .	4

## Problems

4

Clearly, For arbitrary  $a, c \in G$  and  $b, d \in H$

$$\begin{aligned} ac &= ca \wedge bd = db \\ \leftrightarrow (ac, bd) &= (ca, db) \\ \leftrightarrow (a, b)(c, d) &= (c, d)(a, b) \end{aligned}$$

$$\begin{array}{r} 0.75 \\ \hline 0.75 \end{array}$$

I guess the general case is any group-theoretic property on both  $G$  and  $H$  is also on  $G \oplus H$ , and vice versa.

5

Assume for the sake of contradiction  $Z \oplus G$  is cyclic. Then by definition there is a generator  $(a, b)$ . Then necessarily  $\langle a \rangle = Z$  and  $\langle b \rangle = G$  as by definition we have  $(a, b)^k = (a^k, b^k)$ . Observe  $\langle a \rangle$  is of infinite order. Fix  $c \in Z$ , Then we know  $a^k = c$  for some  $k$ . Compute  $(a, b)^k = (a^k, b^k) = (c, d)$ . Let  $h$  be the element other than  $d$  in  $G$ . Now we can't generate  $(c, h)$ . By theorem 4.1 (page 76) if  $a^i = a^k$  then  $i = k$ . In other words,  $k$  is the only integer that yields  $a^k = c$ .

①  
1

for infinite groups

6

Clearly  $(1, 1) \in Z_8 \oplus Z_2$  is of order 8. We claim no element of  $Z_4 \oplus Z_4$  is of order 8, Which suffices to solve the problem.

From Theorem 4.3 (page 81) we know any element of  $Z_4$  is of order, which divides 4. In other words, For any element  $a$ , there is  $k \leq 4$  such that  $k|a| = 4$ . Similarly for another element  $b$  we have  $k'|b| = 4$ .

So for any  $(a, b) \in Z_4 \oplus Z_4$ , Observe  $(a, b)^4 = (a^4, b^4) = (a^{k|a|}, b^{k'|b|}) = ((a^{|a|})^k, (b^{|b|})^{k'}) = (0^k, 0^{k'}) = (0, 0)$ . So order of  $(a, b)$  is at most 4.

$$\begin{array}{r} 0.5 \\ \hline 0.5 \end{array}$$

15

Let  $\phi : \mathbb{C} \rightarrow \mathbb{R} \oplus \mathbb{R}$  where  $\phi(a + bi) = (a, b)$ .

- Injective.  $\phi(a + bi) = \phi(c + di)$  implies  $(a, b) = (c, d)$ , and in turn  $a = c$  and  $b = d$ .  $\Rightarrow a + bi = c + di$ .
- Surjective. For any  $(a, b)$  we have  $\phi(a + bi) = (a, b)$ .
- Preserves Operation.  $\phi(a + bi) + \phi(c + di) = (a, b) + (c, d) = (a + c, b + d) = \phi((a + c) + (b + d)i) = \phi((a + bi) + (c + di))$ .

$$\begin{array}{r} 0.75 \\ \hline 0.75 \end{array}$$

17

Since  $G \oplus H$  is cyclic, it has a generator  $(a, b)$ . It follows  $\langle a \rangle = G$  and  $\langle b \rangle = H$ . If that is not the case, Then we can select an element from  $G$  or  $H$  whereby  $(a, b)^k = (a^k, b^k)$  won't cover it, on it corresponding index.

a.s  
o.s

21

Denote the equivalence  $\langle (g, h) \rangle = \langle g \rangle \oplus \langle h \rangle$  by (1).

Recall theorem 8.1 (page 158).

By definition we know  $(g, h)^k = (g^k, h^k)$  where  $g^k \in \langle g \rangle$  and  $h^k \in \langle h \rangle$ .

The condition is  $|g|$  and  $|h|$  are coprime. Observe it is equivalent to  $lcm(|g|, |h|) = |g||h|$ .

(Necessity) We show given (1), The condition holds. Since sets are equal, and cardinality of L.H.S is  $|g| \cdot |h|$ , Then  $|(g, h)| = |g| \cdot |h|$ . By *thm 8.1*, The condition is satisfied.

(Sufficient) We show given the condition, (1) holds. By *thm 8.1*,  $|(g, h)| = |g| \cdot |h|$ . So its cardinality is the same as R.H.S, and it is a subset of it. It follows (1) holds.

2  
2

23

Any element in  $\mathcal{Z}_3$  is of order 3, except the identity 0. Consider an arbitrary non-identity element  $(x_1, x_2, \dots, x_k) \neq e = \underbrace{(0, \dots, 0)}_{k \text{ times}}$  in  $\underbrace{\mathcal{Z}_3 \oplus \dots \oplus \mathcal{Z}_3}_{k \text{ times}}$ . We claim  $|(x_1, \dots, x_k)| = 3$ .

Following the fact all non-identity elements are of order 3, and we have some  $x_i \neq 0$ ,

$$\begin{aligned} (x_1, x_2, \dots, x_k)^1 &= (x_1^1, x_2^1, \dots, x_k^1) \neq e \\ (x_1, x_2, \dots, x_k)^2 &= (x_1^2, x_2^2, \dots, x_k^2) \neq e \\ (x_1, x_2, \dots, x_k)^3 &= (0, 0, \dots, 0) = e \end{aligned}$$

} Why?  
I can't see any reasoning!

1  
2

Therefore we have  $3^k - 1$  elements of order 3 in  $\underbrace{\mathcal{Z}_3 \oplus \dots \oplus \mathcal{Z}_3}_{k \text{ times}}$ .

The question was about subgroups, not elements!

35

Recall the square root of any complex number  $z$  exists. Observe  $C^*$  is closed under the square root operation.

Assume for the sake of contradiction, there is an isomorphism  $\phi : C^* \rightarrow R^* \oplus R^*$ . Then

by surjectivity there is some complex  $z$  where  $\phi(z) = (-1, -1)$ . It follows


$$\begin{aligned} \phi(\sqrt{z} \cdot \sqrt{z}) &= (-1, -1) \\ \phi(\sqrt{z}) \cdot \phi(\sqrt{z}) &= \\ (\phi(\sqrt{z}))^2 &= \\ (a, b)^2 &= \\ (a^2, b^2) &= \end{aligned}$$



In other words  $a^2 = -1$  and  $b^2 = -1$ , but either of these leads to a contradiction, as no square of a real number is negative.

46

The infinite group is  $\mathcal{Z} \oplus D_4 \oplus A_4$ . Clearly  $\{(e_{\mathcal{Z}}, x, e_{A_4}) \mid x \in D_4\}$  and  $\{(e_{\mathcal{Z}}, e_{D_4}, x) \mid x \in A_4\}$  are both subgroups.



48

**Claim.** It is all permutations on  $\mathcal{Z}_2 \oplus \mathcal{Z}_2$  which maps  $(0, 0)$  to itself.



**Note.** Our characterization is consistent with the fact the identity is always mapped to itself, and that isomorphism is a bijection.

**Fact.** In any group, fixing element  $a_0$ , then for any elements  $b_0 \neq b_1$ , we have  $a_0 b_0 \neq a_0 b_1$ .

**Lemma.** For any  $(a, b) \in \mathcal{Z}_2 \oplus \mathcal{Z}_2$ ,  $(a, b)^2 = (a^2, b^2) = (0, 0) = e$ . As  $0^2 = 0$  and  $1^2 = 0$ .

**Lemma.** Any two elements of  $X = \{(0, 1), (1, 0), (1, 1)\}$  multiplies to the third.

For distinct  $a, b, c \in X$ ,  $ab \neq (0, 0)$  since  $aa = (0, 0)$ . Also  $ab \neq a$  since  $a(0, 0) = a$ . Similarly  $ab \neq b$ . Therefore the only remaining choice is  $ab = c$ .

**Theorem.** Our permutations preserve the operation.

We know for distinct elements  $a, b, c \in X$ , we have  $ab = c$ . As  $\phi$  is a permutation on these, We have  $X = \{\phi(a), \phi(b), \phi(c)\}$ . It follows  $\phi(a)\phi(b) = \phi(c)$ . That concludes  $\phi(c) = \phi(ab) = \phi(a)\phi(b)$ .

